

α spectrum from ^{16}N β decay and the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction

J. Humblet,* B. W. Filippone, and S. E. Koonin

W. K. Kellogg Radiation Laboratory, 106-38, California Institute of Technology, Pasadena, California 91125

(Received 29 April 1993)

New data on the spectrum of α particles following the ^{16}N β decay are used to revise and extend a previous evaluation of the $E1$ part of the astrophysical S factor for the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction.

PACS number(s): 25.90.+k

In a previous calculation [1], we used the measured spectrum [2] of α particles from ^{16}N β decay in an attempt to constrain the astrophysical S factor for the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction. For this test, we used the p -wave part of the total α spectrum that had been separated by Barker [3,4] using an R -matrix parametrization. This test showed the importance of obtaining a better set of data over a much wider energy range, and in particular at lower energies. Here, we update our fit by using a K -matrix parametrization to fit the total α spectrum recently obtained by Buchmann *et al.* [5] at TRIUMF.

The K -matrix parametrization for the total α spectrum reads

$$N_\alpha = N_{1\alpha} + N_{3\alpha}, \quad (1)$$

with, for $l = 1$ and 3 ,

$$N_{l\alpha} = f_\beta(E) p_{l\alpha}^2(E) |n_l|^2 / (1 + d_l^2), \quad (2)$$

and

$$n_l = \sum_{\lambda=1}^3 B_{l\lambda} g_{l\alpha\lambda} / (E_{l\lambda} - E) + b_{l\beta}, \quad (3)$$

$$d_l = p_l^2 K_{l\alpha\alpha}, \quad (4)$$

$$K_{l\alpha\alpha} = \sum_{\lambda=1}^3 g_{l\alpha\lambda}^2 / (E_{l\lambda} - E) + b_{l\alpha\alpha}. \quad (5)$$

The notation is as in Ref. [1], except that we now explicitly include the l label, rather than assuming implicitly $l = 1$ only. The feeding factors B_{11} and B_{31} for the two bound states (1^- at $E_x = 7.12$ MeV, 3^- at $E_x = 6.13$ MeV) are obtained from Eq. (3.16) of Ref. [1] using the branching ratios from Ref. [6]. The parameter $b_{1\beta}$ is set to zero as in Ref. [1]. The calculated α spectrum has been convolved with the detector resolution for comparison with the data, assuming a Gaussian response whose energy independent full width at half maximum is 35 keV.

As in Ref. [1], we minimize

$$\chi_{\text{eff}}^2 = \frac{1}{3} (\chi_{\gamma 1}^2 + \chi_{\delta 1}^2 + \chi_\beta^2), \quad (6)$$

where $\chi_{\gamma 1}^2$, $\chi_{\delta 1}^2$, and χ_β^2 are χ^2 per data point for the $E1$

capture cross section, the p -wave elastic scattering phase shift, and the total α spectrum, respectively.

For the f -wave contribution in Eq. (3), we initially fixed the energies E_{31}, E_{32} at -1.032 MeV and 4.415 MeV, respectively, and fixed the energy (E_{33}) of the background pole term. For the remaining parameters, we considered each of the two products $B_{32}g_{3\alpha 2}, B_{33}g_{3\alpha 3}$ as one single free parameter and included $g_{3\alpha 1}$ and $b_{3\beta}$. However, we noted that the detailed form of the f -wave contribution was only weakly constrained by the data. No additional information could be obtained when the f -wave elastic scattering phase shift δ_3 was simultaneously fitted. Apparently, the 3^- resonance at $E_{32} = 4.415$ MeV is too far above the energy range ($E = 1.3 - 1.5$ MeV) that determines the f -wave part of the α spectrum to have a significant effect. After many trials, we concluded that χ_{eff}^2 could not be improved by using more than one free parameter for n_3 .

The best result is obtained with a constant n_3 ,

$$n_3 = b_{3\beta}, \quad (7a)$$

but choosing

$$n_3 = B_{31}g_{3\alpha 1} / (E_{31} - E) \quad (7b)$$

gives only a small increase of χ_{eff}^2 . The latter form for n_3 is the one adopted in Ref. [5].

The $l = 3$ phase shift also plays a role in determining the denominator of $N_{3\alpha}$ through d_3^2 . However, in a separate best fit of the δ_3 phase shift, d_3^2 remains much smaller than 1 in the energy range of interest. At $E = 1.4, 2.4$, and 3.2 MeV, d_3^2 is 9.4×10^{-9} , 1.1×10^{-4} , and 1.0×10^{-2} , respectively. Accordingly, we have not varied the parameters of $K_{3\alpha\alpha}$ in d_3 in order to reduce the number of free parameters and better constrain them.

The data we used for the δ_1 phase shift and the $E1$ capture cross section are the same as in Refs. [1,7]. From the TRIUMF data, we removed four data points [8] ($E_\alpha = 1.99 - 2.05$ MeV) at energies close to the 2^+ resonance at $E = 2.68$ MeV. This is important for the α spectrum as there is a known β -decay branching ratio to this state ($\simeq 7 \times 10^{-9}$), and we have not included this state in our parametrization.

The results of two fits corresponding to Eqs. (7a) and (7b), respectively, are given in Tables I and II, while Figs. 1–4 correspond to Eq. (7a) only. The first fit is slightly better than the second one, due to the apparently better parametrization of the f -wave part of the α spec-

*Permanent address: Université de Liège, Institut de Physique B5, Sart Tilman, B-4000 Liège 1, Belgium.

TABLE I. Parameter values for two best fits corresponding to n_3 given by Eq. (7a) (second column) and by Eq. (7b) (third column). As in Ref. [1], $a = 5.46$ fm, and E_{13} is an echo pole. The ranges for S_{E1} correspond approximately to one standard deviation.

E_{11} (MeV)	(-0.0451)	(-0.0451)
$g_{1\alpha 1} a^{-3/2}$ (MeV $^{1/2}$)	-5.408	-5.214
$g_{1\gamma 1} a^{-3/2}$ (MeV $^{1/2}$)	(1.897×10^{-3})	(1.897×10^{-3})
B_{11}	(1193)	(1193)
E_{12} (MeV)	2.4435	2.4437
$g_{1\alpha 2} a^{-3/2}$ (MeV $^{1/2}$)	7.007	7.030
$g_{1\gamma 2} a^{-3/2}$ (MeV $^{1/2}$)	0.6555×10^{-3}	0.6572×10^{-3}
B_{12}	-410.5	-412.8
E_{13} (MeV)	(8.000)	(8.000)
$g_{1\alpha 3} a^{-3/2}$ (MeV $^{1/2}$)	14.53 i	14.92 i
$g_{1\gamma 3} a^{-3/2}$ (MeV $^{1/2}$)	$-2.550 \times 10^{-3} i$	$-2.641 \times 10^{-3} i$
B_{13}	700.5 i	644.9 i
$b_{1\alpha\alpha} a^{-3}$	77.72	77.95
$b_{1\alpha\gamma} a^{-3}$	-5.797×10^{-3}	-5.887×10^{-3}
E_{31} (MeV)	(-1.032)	(-1.032)
$g_{3\alpha 1} a^{-7/2}$ (MeV $^{1/2}$)	—	6.780×10^{-2}
B_{31}	—	(2458)
$b_{3\beta} a^{-7}$	-66.75	—
$\Gamma_{1\gamma 1}$ (MeV)	(55×10^{-9})	(55×10^{-9})
$\Gamma_{1\alpha 2}$ (MeV)	0.454	0.457
$\Gamma_{1\gamma 2}$ (MeV)	16×10^{-9}	16×10^{-9}
S_{E1} (MeV b) at χ^2_{\min}	0.045	0.042
S_{E1} (MeV b) range	0.039 – 0.050	0.035 – 0.046

trum with n_3 given by Eq. (7a) rather than by Eq. (7b). The $\chi^2_{\gamma 1}$, $\chi^2_{\delta 1}$ are practically the same for both fits and for our best fit of 1991 [1] using the Wäffler data. Our present results for χ^2_{β} are significantly larger than those we obtained in Ref. [1]. The uncertainties assigned to the p -wave α data used in Ref. [1] appear to be dominated by a nonstatistical uncertainty in the α particle energy resulting in the very low χ^2_{β} we obtained previously.

For two different fits to the α spectrum, Ref. [5] has obtained $\chi^2_{\beta} = 1.46$ and 1.30. While these are somewhat better than ours, Ref. [5] did not simultaneously fit the $l = 1$ phase shift. Not fitting δ_1 has two other consequences. First, the energy of the 1^- resonance is lowered to $E_{12} \approx 2.41$ MeV and the corresponding partial width is reduced to $\Gamma_{1\alpha 2} = 0.371$ MeV. Second, fitting δ_1 data such as those of Plaga *et al.* [9] constrains $g_{1\alpha 1}$ to lower values, explaining our lower S factor compared to Ref. [5]. We do not think one can disregard fitting the presently available data for δ_1 , but this problem and the above remarks certainly require further study. More accurate and extended data for δ_1 at low energies would be desirable,

TABLE II. Values of χ^2 for simultaneous fits to the three sets of data and the minimum values of χ^2_{eff} .

	$\chi^2_{\gamma 1}$	$\chi^2_{\delta 1}$	χ^2_{β}	χ^2_{\min}
In Ref. [1]	0.88	1.51	0.20	0.86
Fit 1	0.87	1.51	1.62	1.33
Fit 2	0.88	1.51	1.78	1.39




FIG. 1. Fit to the $E1$ S -factor data of Ref. [10] when the δ_1 phase shift and the α spectrum N_α are fitted simultaneously.




FIG. 2. Fit to the same three sets of phase shift data used in Ref. [1] when the $E1$ S factor and α spectrum N_α are fitted simultaneously.




FIG. 3. Fit of the α spectrum to the data of Ref. [5] when the $E1$ S factor and the δ_1 phase shift are simultaneously fitted. The solid curve, the dashed curve, and the dot-dashed curve are the fits of N_α , $N_{1\alpha}$, and $N_{3\alpha}$, respectively.




FIG. 4. The effective χ^2 vs $S_{E1}(0.3)$. The minimum value of χ^2_{eff} is obtained for a series of fixed values assigned to $S_{E1}(0.3)$, in the range 0.00–0.12 MeV b. The solid curve corresponds to the first fit in Table I, and the dotted one, shown for comparison, corresponds to the first fit in Ref. [1]. From such curves, we extract the range of allowed values for $S_{E1}(0.3)$.

as well as a better determination of $B_{11} = 1193 \pm 54$. This feeding factor is strongly correlated with $g_{1\alpha 1}$ in $N_{1\alpha}$. It is deduced from the branching ratios given in Ref. [6], and its error is dominated by the branching ratio to the 7.12 MeV bound state.

For both fits reported in Table I, the term involving E_{13} is an echo pole. This implies $g_{1\alpha 3}^2 < 0$; i.e., an imaginary $g_{1\alpha 3}$, so that $g_{1\gamma 3}$ and B_{13} must also be imaginary.

An echo pole is necessary to parametrize the downward trend of δ_1 at $E > 3.1$ MeV, but no restriction is imposed on the sign of the real products $g_{1\alpha 3}g_{1\gamma 3}$, $B_{13}g_{1\alpha 3}$. (In an R -matrix fit, a downward trend in the total phase shift is insured by its hard-sphere contribution.)

The χ^2 we have obtained for the capture data, $\chi^2_{\gamma 1} = 0.87$ and 0.88 , are much lower than those reported in Table I of Ref. [5], corresponding to $\chi^2_{\gamma 1} = 2.73$ and 2.69 . This is obviously related to the fact that we have fitted $S_{E1}(E)$ only to the data of Kremer *et al.* [10], as in Ref. [1], while in Ref. [5] four sets of independent data are simultaneously fitted.

From the results in Table I and Fig. 4, we obtain a best value for $S_{E1}(0.3)$ of

$$S_{E1}(0.3) = 0.045^{+0.005}_{-0.006} \text{ MeV b}, \quad (8)$$

where the errors correspond approximately to one standard deviation (increase in χ^2_{eff} of 5%, see Ref. [11]). For comparison with Ref. [1], where we quote 95% confidence level errors bars, our errors would be $^{+0.010}_{-0.012}$ (increase in χ^2_{eff} of 20%).

The authors are grateful to R. E. Azuma, C. A. Barnes, and L. Buchmann for an early communication of their data and details about their fits. J.H. would like to thank his colleagues, G. Cantraine and J. Destin  , from the University of Li  ge, for the use of their computing facilities, and G. de Brus for his assistance. This work was supported in part by the National Science Foundation, Grant Nos. PHY90-13248 and PHY91-15574.

- [1] J. Humblet, B. W. Filippone, and S. E. Koonin, Phys. Rev. C **44**, 2530 (1991).
- [2] H. H  ttig, K. H  nchen, and H. W  ffler, Phys. Rev. Lett. **25**, 941 (1970); K. Neubeck, H. Schober, and H. W  ffler, Phys. Rev. C **10**, 320 (1974).
- [3] F. C. Barker, Aust. J. Phys. **24**, 777 (1971).
- [4] F. C. Barker, private communication.
- [5] L. Buchmann, R. E. Azuma, C. A. Barnes, J. M. D'Auria, M. Domskey, U. Giesen, K. P. Jackson, J. D. King, R. G. Korteling, P. McNeely, J. Powell, G. Roy, J. Vincent, T. R. Wang, S. S. M. Wong, and P. R. Wrean, Phys. Rev. Lett. **70**, 726 (1993).
- [6] F. Ajzenberg-Selove, Nucl. Phys. **A460**, 1 (1986), see p.

50.

- [7] B. W. Filippone, J. Humblet, and K. Langanke, Phys. Rev. C **40**, 515 (1989).
- [8] These four points have not been removed from our Fig. 3.
- [9] R. Plaga, H. W. Becker, A. Redder, C. Rolfs, H. P. Trautvetter, and K. Langanke, Nucl. Phys. **A465**, 291 (1987).
- [10] R. M. Kremer, C. A. Barnes, K. H. Chang, H. C. Evans, B. W. Filippone, K. H. Hahn, and L. W. Mitchell, Phys. Rev. Lett. **60**, 1475 (1988).
- [11] Particle Data Group, K. Hikasa *et al.*, Phys. Rev. D **45**, III. 32 (1992).